



<https://bjm.ui.ac.ir/?lang=en>

Journal of Microbial biology  
E-ISSN: 2322-5173  
12<sup>th</sup> Year, Vol. 12, No. 48, Winter 2023 pp. 1-16  
Received: 06-06-2023 Accepted: 03-08-2023

**(Research Paper)**

## **Enhanced Nitrogen Removal by a Heterotrophic Nitrifying-Aerobic Denitrifying Bacterial Consortium with Potential for the Treatment of High-strength Ammonium Wastewater**

**Noushan Hatami**

Department of Cell and Molecular Biology & Microbiology, Faculty of Biological Science and Technology, University of Isfahan, Isfahan, Iran, nooshanhatami@yahoo.com

**Zahra Etemadifar\*** 

Department of Cell and Molecular Biology & Microbiology, Faculty of Biological Science and Technology, University of Isfahan, Isfahan, Iran, z.etemadifar@sci.ui.ac.ir

**Rasoul Shafiei**

Department of Cell and Molecular Biology & Microbiology, Faculty of Biological Science and Technology, University of Isfahan, Isfahan, Iran, r.shafiei@sci.ui.ac.ir

### **Abstract**

**Introduction:** Nitrogen compounds are used in various industries and cause pollution of water and soil. As a result, the bioremediation of nitrogen compounds from the environment becomes an important issue. In this way, the use of microorganisms as a consortium has enhanced bioremediation. This study aimed to remove inorganic nitrogen forms by heterotrophic nitrifying-aerobic denitrifying (HN-AD) bacterial consortium and compare it to pure cultures.

**Materials and Methods:** HN-AD bacteria were isolated from soil and aquarium water and identified. After that, the bacterial consortium was prepared from two isolated bacteria and two strains from the microbiological culture collection of the University of Isfahan (Iran). The ability to remove ammonium, nitrite, and nitrate was assayed using spectrophotometry for both pure cultures and the consortium.

---

\* Corresponding Author  
2322-5181/ © 2023 The Authors

This is an open access article under the CC-BY-NC-ND 4.0 License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)



**Results:** *Pseudomonas mendocina* AquaN and *Pseudomonas monteilii* Nht were isolated and identified for their efficient capabilities in HN-AD. The bacterial consortium including two isolated bacteria and *Acinetobacter calcoaceticus* SCC2, *Rhodococcus erythropolis* R1 was able to consume  $\text{NH}_4^+\text{-N}$  (85.6  $\text{mg.L}^{-1}$ ),  $\text{NO}_2^-\text{-N}$  (23.1  $\text{mg.L}^{-1}$ ), and  $\text{NO}_3^-\text{-N}$  (102.5  $\text{mg.L}^{-1}$ ) as sole nitrogen sources by removal efficiencies 98.4%, 90.5%, and 40.0%, respectively that were better than pure cultures performances. Specifically, the consortium was capable of removing 250 to 2045  $\text{mg.L}^{-1}$   $\text{NH}_4^+\text{-N}$  with up to 95% efficiency within 48h.

**Discussion and Conclusion:** The compatibility of the strains as a consortium was considerable and the bacterial consortium showed significant performances in nitrogen removal, especially the removal of 2045  $\text{mg.L}^{-1}$   $\text{NH}_4^+\text{-N}$ . These findings indicated the bacterial consortium could be a promising candidate for the treatment of high-strength ammonium wastewater.

**Keywords:** *Acinetobacter*, Consortium, Heterotrophic nitrification, High-strength ammonium removal, *Pseudomonas*, *Rhodococcus*.

## Introduction

Nitrogen compounds are used in a variety of industrial and agricultural processes. Due to their wide use, they are frequently found in soil, air, wastewater, and aquatic environments (1). Inorganic nitrogen is present in ammonium, nitrite, and nitrate forms. Ammonium lowers the quality of drinking water, and makes a serious danger to human health because of eutrophication, is toxic to aquaculture, and changes the structure of microbial communities (2, 3). High nitrate concentrations can cause methemoglobinemia or blue baby syndrome in newborns. In addition, nitrate is reduced to nitrite and carcinogenic chemicals nitrosamines in the digestive system, which cause several forms of cancer, including gastric cancer (4, 5).

The concentration of ammonium nitrogen is 1-10  $\text{mg.L}^{-1}$  in the soil without fertilizer, 50  $\text{mg.L}^{-1}$  in areas contaminated with agricultural fertilizers and 1000  $\text{mg.L}^{-1}$  in industrial wastewater (6). In sum, due to the concentration of ammonium in soil, and all the existing problems need to be paid attention to nitrogen removal in wastewater.

Nitrogen forms in the water and

wastewater can be removed by physicochemical and biological methods. Physicochemical treatments include reverse osmosis, ion exchange, electrodialysis, and activated carbon technology systems (7). Biological methods can perform in the conventional way that autotrophic bacteria are involved. Conventionally, ammonium is oxidized to nitrite and nitrate under aerobic conditions by autotrophic nitrifying bacteria. Nitrate could be transformed to nitrogen gaseous by heterotrophic nitrate-reducing bacteria under anaerobic conditions (8). This method is time-consuming and small variations in temperature, pH, and substrate concentration affect the activity and growth of autotrophic nitrifiers (9, 10).

During the past few years, a new method, heterotrophic nitrification-aerobic denitrification (HN-AD) has been studied. HN-AD is done by heterotrophic bacteria such as *Pseudomonas* (8), *Acinetobacter* (11), *Alcaligenes* (12), *Klebsiella* (13), and *Bacillus* (14). In this manner, ammonium can be converted into gaseous nitrogen under aerobic conditions in one reactor (12). HN-AD bacteria generally remove nitrogen by

using carbon sources like sodium citrate, sodium acetate, glucose, and sodium succinate (9). In comparison to the conventional method, HN-AD exhibited a fast-growing rate, simultaneous ammonium and nitrate or nitrite removal, less contamination, and cost-effectiveness (15). There are different biochemical pathways of HN-AD in bacteria (16).

Most studies about HN-AD were done in pure culture of bacteria; less attention has been paid to co-culture or consortium of HN-AD bacteria. The application of microbial consortium is beneficial over pure culture because synergistic interactions between different strains can provide the use of complex carbon sources, increase each other's growth by cross-feeding, managing their environment, and enhancing productivity (17). The consortium of HN-AD bacteria with various nitrogen removal pathways may be more efficient than one pure culture in complex environments (7).

This study focused on the HN-AD consortium with high nitrogen removal efficiency. The specific goals of the present study were 1) isolation of HN-AD bacteria, 2) assay strains interactions, 3) determination of the cell growth and nitrogen removal characteristics of pure cultures and bacterial consortium, and 4) assay high-strength ammonium removal by HN-AD bacterial consortium.

## Materials and Methods

**Bacterial strains:** The HN-AD bacteria were isolated from the soil of the garden area at the University of Isfahan and the aquarium freshwater of the aquaculture center in Isfahan, Iran. In addition, two strains from the microbiological culture collection of the University of Isfahan (Iran) *Rhodococcus erythropolis* R1 (GU570564.1) (18) and *Acinetobacter calcoaceticus* SCC2 (KU685396.1) (19) were used in this research.

**Media and Chemicals:** Modified heterotrophic nitrification medium (HNM) contained (g.L<sup>-1</sup> of distilled water): (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> 0.33, sodium succinate 2.8, KH<sub>2</sub>PO<sub>4</sub> 0.5, Na<sub>2</sub>HPO<sub>4</sub> 0.5, MgSO<sub>4</sub>.7H<sub>2</sub>O 0.2, NaCl 3.0, trace element solution 2.0 mL, and pH 7 ± 0.2 (26). Nitrite denitrification medium (DM-1) and nitrate denitrification medium (DM-2) are based on HNM but DM-1 consisted of NaNO<sub>2</sub> (0.05 g.L<sup>-1</sup>) and DM-2 contained KNO<sub>3</sub> (0.722 g.L<sup>-1</sup>) instead of (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> as a nitrogen source. The trace element solution contains (g.L<sup>-1</sup> of distilled water) EDTA 57.1, ZnSO<sub>4</sub> 3.9, CaCl<sub>2</sub> 7, FeSO<sub>4</sub>.4H<sub>2</sub>O 5, (NH<sub>4</sub>)<sub>6</sub>Mo<sub>7</sub>O<sub>24</sub>.4H<sub>2</sub>O 1.1, CoCl<sub>2</sub>.6H<sub>2</sub>O 1.6, and CuSO<sub>4</sub>.5H<sub>2</sub>O 1.6 (20). The Luria Bertani (LB) broth contained (g.L<sup>-1</sup>) peptone 10.0, yeast extract 5.0, and NaCl 10.0. Solid media were prepared from mentioned media with an addition of 1.5% agar. The media were sterilized by autoclave for 15 min at 121°C. Nessler's reagents were purchased as Merck nessler's reagents 109011 and 109028 from Merck Germany. Griess's reagent was purchased as Griess's reagent 03553 from Sigma-Aldrich USA. All the chemicals used were of analytical grade.

**Isolation of HN-AD Bacteria:** To enrich heterotrophic nitrification-aerobic denitrification bacteria, at first, the soil sample (10 g) was added to 100 mL Ringer's solution and then shaken at room temperature. Secondly, 5 mL of soil solution was mixed with 95 mL HNM and incubated at 30°C on a rotary shaker at 120 rpm (modified (21)). Aquarium water samples (50 mL) were added to 250 mL conical flasks containing 50 mL sterile liquid HNM and incubated at 30°C on a rotary shaker at 120 rpm. After three days, 5 mL of the cell suspension from each culture was transferred to 95 mL fresh HNM and incubated at 30°C and 120 rpm for three days. The enriched bacterial cultures were gradient-diluted, and 0.1 mL of each diluted

medium was spread on solidified HNM medium and incubated at 30°C for two days under aerobic conditions. Each separate colony was isolated and purified. The nitrification ability of the isolates was determined in HNM broth and the isolates with the maximum nitrification ability were selected (8).

#### **Molecular and Biochemical Identification:**

The best two isolates were identified by morphological, biochemical, and molecular methods. Gram staining and biochemical tests include catalase, SH<sub>2</sub> production, indole, anaerobic nitrate reduction, V-P test, methyl red test, urease, oxidase, hydrolysis of starch, O/F test, using citrate, production of acid from glucose, fructose, lactose, and mannitol (22).

16S rRNA genes of the selected isolates were amplified by PCR for molecular identification. The DNA was extracted from bacterial suspension by a DNA extraction kit (Cinnagen Co., Iran). The universal primers 27F (5'-AGAGTTTGATCMTGGCTCAG-3') and 1492R (5'-TACGGYTACCTTGTACGACTT-3') were used (23). The PCR program was 35 cycles and steps were carried out as follows: denaturation at 95°C for 5 min, 94°C for 45 seconds, annealing at 55°C for 1 min, extension at 72°C for 1 min; and final extension at 72°C for 7 min. The PCR products were analyzed by electrophoresis on the 1% agarose gel and sequenced by the Sanger method (Model ABI 3730 XL, FAZA Biotech Co., Iran). To compare sequences with other microorganisms, sequence alignment was applied by using the Basic Local Alignment Search Tool (BLAST; <http://www.ncbi.nlm.nih.gov/BLAST/Blast.cgi>). The 16S rRNA gene sequences were submitted to the GenBank database. A neighbor-joining tree was created using the MEGA 6.0 program.

#### **Inoculum Preparation and Culture**

**Conditions:** To prepare the inoculum, a single pure colony of each strain was transferred to 20 mL of fresh media including HNM, DM-1, and DM-2 as pre-cultures. All media were incubated at 30°C with shaking at 120 rpm to cells reached the middle of the logarithmic phase and the cell concentration became equal to 0.5 McFarland. Afterward, 5 % (v/v) of inoculum was transferred to an Erlenmeyer flask containing 95 ml of HNM, DM-1, and DM-2 broth media. To make consortium inoculation, inoculum solutions of strains were mixed in equal volumes and the total inoculum size reached 5% (v/v).

#### **Heterotrophic Nitrification/Aerobic Denitrification Capability of Pure Cultures:**

To assess the capabilities of ammonium, nitrite, and nitrate removal, 5% (v/v) of each pure culture suspension was inoculated into HNM (100 mg.L<sup>-1</sup> of NH<sub>4</sub><sup>+</sup>-N), DM-1 (10 mg.L<sup>-1</sup> NO<sub>2</sub><sup>-</sup>-N), and DM-2 (110 mg.L<sup>-1</sup> of NO<sub>3</sub><sup>-</sup>-N), respectively. The flasks were incubated at 30°C and 120 rpm under aerobic conditions. A medium without inoculation was used as a control for each examination. The samples were periodically taken to determine the concentration of NH<sub>4</sub><sup>+</sup>-N, NO<sub>2</sub><sup>-</sup>-N, NO<sub>3</sub><sup>-</sup>-N, and optical density at 600 nm (OD<sub>600</sub>). After 24 h incubation, total nitrogen (TN) was measured in all media.

#### **Bacterial Consortium Design**

**Consortium Pre-Tests:** To find out the interactions between the strains, two tests were done. The first test was the Bilayer culture that one strain was cultivated as a circular form on the nutrient agar plate center, and then the second strain was cultivated by the pour plate method on the top layer (24). The cross streak method was used as the second test that one strain was cultivated as a linear form on the center of HNM agar and the other strains were cultivated vertically by 1 cm space from the line of the culture (modified (25)).

### Assessment of Nitrogen Removal Capability of Bacterial Consortium

**Assessment of Nitrogen Removal Capability of Bacterial Consortium in the Growth Phase:** To determine the performance of nitrogen removal by the bacterial consortium in the growth phase, strains were inoculated to HNM, DM-1, and DM-2. Samples were taken regularly to determine  $\text{NH}_4^+\text{-N}$ ,  $\text{NO}_2^-\text{-N}$ ,  $\text{NO}_3^-\text{-N}$ , and  $\text{OD}_{600}$ . Media without inoculation were used as controls.

**Assessment of Ammonium Removal by Resting Cells of Bacterial Consortium:** For the preparation of consortium resting cells, 1% (v/v) inoculum of 0.5 McFarland concentration of each strain was inoculated to 50 ml of LB Broth in 250 ml Erlenmeyer flasks and incubated at 30°C and 120 rpm agitation. At the end of the exponential growth phase, cells were harvested by centrifugation (Z 300 k, Hermle Labortechnik, Germany) at 6000 rpm and 14°C for 6 min. The pellets were washed twice with sodium phosphate buffer (pH 7.2). Afterward, prepared resting cells of all strains were resuspended in 100 mL of HNM. The  $\text{OD}_{600}$  of the resting cell suspension was adjusted to 2.3. After incubation at 30°C and 120 rpm, samples regularly were taken at 2 h intervals to evaluate  $\text{NH}_4^+\text{-N}$ ,  $\text{NO}_2^-\text{-N}$ , and  $\text{NO}_3^-\text{-N}$ . Total nitrogen (TN) was measured at the end of the incubation period.

**High-strength Ammonium Removal Performance by Bacterial Consortium:** To measure the high-strength ammonium removal efficiency of the bacterial consortium, the initial  $\text{NH}_4^+\text{-N}$  concentration was 250, 425, 575, 740, 990, 1150, 1650, and 2045  $\text{mg.L}^{-1}$ . All the experiments were carried out with 5% (v/v) inoculation, and then the flasks were incubated at 30°C and 120 rpm agitation.

Nitrification rate and efficiency were calculated after 24 and 48 h incubation. The nitrification capability of each strain in the medium contained 2045  $\text{mg.L}^{-1}$  of  $\text{NH}_4^+\text{-N}$  was also assayed for comparison with the consortium.

**Analytical Methods:** To measure some parameters, the samples were centrifuged (Sigma 1-14, Sigma Laborzentrifugen, Germany) at 11000 rpm for 4 min and filtered if required. Levels of total nitrogen and  $\text{NO}_3^-\text{-N}$  were measured according to the standard methods (26); briefly, total nitrogen was assayed based on the persulphate digestion method.  $\text{NO}_3^-\text{-N}$  was calculated by the absorbance at 220 nm and subtracting two times the absorbance at 275 nm.  $\text{NH}_4^+\text{-N}$  was determined by Nessler's reagent spectrophotometry at a wavelength of 420 nm (27).  $\text{NO}_2^-\text{-N}$  was determined by Griess's reagent spectrophotometry at a wavelength of 540 nm (28). The pH was measured by a pH meter (TS-Technology, Iran). The cell growth was determined by spectrophotometry  $\text{OD}_{600}$  (PD-303, Apel, Japan).

**Statistical Analysis:** The data were presented as means  $\pm$  standard deviation (SD) of three replicates. Nitrogen removal rate (N.R.R.) and nitrogen removal efficiency (N.R.E.) were calculated using the following formulas, respectively:

$$\text{N.R.R.} = (C_0 - C_t) / t.$$

$$\text{N.R.E. (\%)} = 100 \times (C_0 - C_t) / C_0$$

$C_0$  is the initial concentration ( $\text{mg.L}^{-1}$ ) of nitrogen,  $C_t$  ( $\text{mg.L}^{-1}$ ) is the final concentration at a given time, and  $t$  is the time (h) that the concentration of nitrogen becomes constant (29).

The statistical analysis was performed by one-way ANOVA with Duncan's test ( $P < 0.05$ ) in SPSS Statistics 22.0 software. Graphical works were carried out by Microsoft Excel 2016.

## Results

### Isolation and Identification of Isolates:

After enrichment and screening, two strains were selected based on heterotrophic nitrification and aerobic denitrification performances. Nht isolate was a Gram-negative bacterium with rod-shaped, its colony on nutrient agar was cream in color, rounded with a regular edge, convex, and sticky. AquaN isolate was a Gram-negative rod-shaped bacterium and its colonies on nutrient agar were light pink, rounded with a regular edge, convex, and opaque. Both the Nht and AquaN isolates showed negative results for the production of SH<sub>2</sub>, indole, anaerobic nitrate reduction, V-P test, methyl red test, and acid production from fructose, lactose, and mannitol. Both the Nht and AquaN isolates showed positive results for catalase, oxidase, urease, motility, and using citrate. The Nht isolate was positive in the acid production from glucose and it was negative for the O/F test. The AquaN isolate showed negative results for the production of acid from glucose and O<sup>+</sup>/F<sup>-</sup>.

The amplified 16S rRNA gene of Nht and AquaN isolates were shown in Figure 1. The BLAST results indicated that the Nht isolate was closely related to *Pseudomonas monteilii*, *P. taiwanensis*, *P. putida*, and *P. mosselii* with 99.71% similarity. Finally, the Nht isolate was identified as *Pseudomonas monteilii* by biochemical tests (Table 1). Also, the AquaN isolate was closely related to *Pseudomonas mendocina*, with a similarity of 99.29%. The partial 16S rRNA gene sequence data of Nht and AquaN strains have been deposited in “GenBank” with the accession numbers MN167837.1 and MN435149.1, respectively. The constructed phylogenetic trees of Nht and AquaN strains based on the 16S rRNA gene sequence of the isolates and some other phylogenetically related strains by the neighbor-joining method were shown in Figure 1. The topology of trees, supported by high

bootstrap values, clearly showed that strain AquaN is within the *Pseudomonas mendocina*, and strain Nht is most probably similar to *Pseudomonas monteilii*.

### The Nitrogen Removal Performance of Pure Cultures

**Pure Cultures Nitrification Performances:** The ammonium removal capability and the growth of four strains include *Pseudomonas monteilii* Nht, *Acinetobacter calcoaceticus* SCC2, *Pseudomonas mendocina* AquaN, and *Rhodococcus erythropolis* R1 were depicted in Figure 2. NH<sub>4</sub><sup>+</sup>-N removal by the Nht strain was started after the lag phase (6 h) and completely removed at 9 h, but the cell growth was continued up to 18 h (Fig. 2a). Nitrite and nitrate were not produced by the Nht strain in this process and total nitrogen was reduced by 47.7%. The strain SCC2 removed NH<sub>4</sub><sup>+</sup>-N during the log phase of cell growth after the short lag phase (3 h), and complete ammonium removal occurred at 9 h but cell growth continued to 21 h (Fig. 2b). No nitrite was produced during nitrification and just 5 mg.L<sup>-1</sup> nitrate was produced and total nitrogen was removed by 31.2% in this process. The AquaN strain removed NH<sub>4</sub><sup>+</sup>-N in the log phase after 6 h start-up time (Fig. 2c). This strain completely degraded ammonium after 12 h, while cell growth continued up to 15 h. During this process, the AquaN strain did not produce nitrite and just 5 mg.L<sup>-1</sup> nitrate was produced at 15 h and then decreased. Total nitrogen was reduced by the AquaN strain by 30.7%. R1 strain was a slow-growing bacterium and its growth phase started after 6 h and removed ammonium from 6 h to 36 h and cell growth was stopped by ammonium ending (Fig. 2d). Nitrite was not produced during nitrification and just 4 mg.L<sup>-1</sup> nitrate was produced. No total nitrogen depletion was observed in the R1 strain after 24 h.




Fig. 1- The phylogenetic tree derived from the neighbor-joining analysis of partial 16S rRNA gene sequences. Bootstrap values (%) are based on 1000 replicates. Bar 0.01 substitutions per nucleotide position. (a,b). PCR amplification results for 16S rRNA genes (M: 10kb DNA Ladder; 1: *Pseudomonas mendocina* strain AquaN; 2: *Pseudomonas monteilii* strain Nht) (c).

Table 1- Differential Biochemical Characteristics of the Nht Isolate from Related *Pseudomonas* Type Strains

| Bacterium                          | Growth on 6% NaCl | Mannitol utilizing | Growth at 45 °C | Growth at 4 °C |
|------------------------------------|-------------------|--------------------|-----------------|----------------|
| Isolate Nht                        | +                 | -                  | -               | -              |
| <i>Pseudomonas monteilii</i>       | +                 | -                  | -               | -              |
| <i>Pseudomonas taiwanensis</i>     | +                 | -                  | +               | +              |
| <i>Pseudomonas mosselii</i>        | +                 | +                  | -               | -              |
| <i>Pseudomonas putida</i>          | -                 | -                  | -               | +              |
| <i>Pseudomonas plecoglossicida</i> | -                 | -                  | -               | -              |




Fig. 2- Ammonium removal characteristics of pure cultures in HNM at 30 °C. The inoculum size was 5% v/v. a: *P. monteilii* Nht, b: *A. calcoaceticus* SCC2, c: *P. mendocina* AquaN d: *R. erythropolis* R1. Error bars indicate the standard deviation of the mean for three replicates. Figures characterize NH<sub>4</sub><sup>+</sup>-N (diamond), NO<sub>2</sub><sup>-</sup>-N (circle), NO<sub>3</sub><sup>-</sup>-N (square), and OD<sub>600</sub> (triangle)

**Aerobic Denitrification Performance by Pure Cultures:** Aerobic denitrification characteristics of Nht, SCC2, AquaN, and R1 strains were investigated in DM-1 and DM-2 for nitrite (Fig. 3) and nitrate (Fig. 4), respectively. The maximum NO<sub>2</sub><sup>-</sup>-N concentration removal by the strains was obtained at 10 mg.L<sup>-1</sup>. Nht, SCC2, and AquaN strains removed NO<sub>2</sub><sup>-</sup>-N at the growth phase for 9 h. The R1 strain started nitrite removal in the stationary phase after 12 h incubation. In all strains, when nitrite started to decrease, nitrate increased. There was a small amount of nitrate at the beginning of the tests which was probably due to the chemical oxidation of nitrite (30).

Strains had shown different behaviors to remove nitrate. Nht, SCC2, and AquaN strains grew in DM-2 and their OD<sub>600</sub> reached 0.6 after 24 h. Due to the slow growth of the R1 strain, nitrate removal lasted up to 72 hours and its OD<sub>600</sub> reached 0.44. SCC2, R1, and AquaN showed increasing NO<sub>2</sub><sup>-</sup>-N concentration at the same time as NO<sub>3</sub><sup>-</sup>-N reduction. In the performances by Nht, SCC2, and AquaN strains, after nitrate decreasing at 24 h, a step of nitrate enhancement was observed simultaneously with the stationary phase. This may be due to the disruption of the cells and the release of nitrate inside them. The Nht strain did not produce NO<sub>2</sub><sup>-</sup>-N during denitrification.




Fig. 3- Nitrite removal characteristics of pure cultures in DM-1 at 30 °C. The inoculum size was 5% v/v. a: *P. monteilii* Nht, b: *A. calcoaceticus* SCC2, c: *P. mendocina* AquaN d: *R. erythropolis* R1. Error bars indicate the standard deviation of the mean for three replicates. Figures characterize  $\text{NO}_2^-$ -N (circle),  $\text{NO}_3^-$ -N (square), and  $\text{OD}_{600}$  (triangle)




Fig. 4- Nitrate removal characteristics of pure cultures in DM-2 at 30 °C. The inoculum size was 5% v/v. a: *P. monteilii* Nht, b: *A. calcoaceticus* SCC2, c: *P. mendocina* AquaN d: *R. erythropolis* R1. Error bars indicate the standard deviation of the mean for three replicates. Figures characterize  $\text{NO}_3^-$ -N (square),  $\text{NO}_2^-$ -N (circle), and  $\text{OD}_{600}$  (triangle)

**Nitrogen Removal Performance by Bacterial Consortium:** In the beginning, two interaction tests were done. In the bilayer test, the effect of one strain on another was assayed on nutrient agar which showed no inhibition zone for all strains. Also, an inhibition effect was not observed by the cross-streak method on HNM. The bacterial consortium was made of *A. calcoaceticus* SCC2, *R. erythropolis* R1, *P. mendocina* AquaN, and *P. monteilii* Nht.  $\text{NH}_4^+\text{-N}$  was removed as  $85.3 \text{ mg.L}^{-1}$  by the bacterial consortium during 9 h, while at this time, nitrate and nitrite were not produced (Fig. 5a). The average ammonium removal rate

and efficiency of the bacterial consortium and pure cultures were shown in Table 2.

The aerobic denitrification characteristics of the bacterial consortium were assayed in DM-1 and DM-2 as depicted in Figures 5b and c. In contrast to pure cultures, the bacterial consortium was able to degrade  $23 \text{ mg.L}^{-1} \text{ NO}_2^-\text{-N}$  and began nitrite removal as soon as it was inoculated and the growth was stopped when nitrite was completely consumed. During nitrite removal,  $\text{NO}_3^-\text{-N}$  concentration increased within 12 h and the final concentration of nitrate was  $22.5 \text{ mg.L}^{-1}$ .




Fig. 5- Nitrogen removal characteristics by the bacterial consortium at 30 °C. The inoculum size was 5% v/v. a: Ammonium removal characteristics by growing cells in HNM, b: Nitrite removal characteristics in DM-1, c: Nitrate removal characteristics DM-2. Error bars indicate the standard deviation of the mean for three replicates. Figures characterize  $\text{NH}_4^+\text{-N}$  (diamond),  $\text{NO}_2^-\text{-N}$  (circle),  $\text{NO}_3^-\text{-N}$  (square), and  $\text{OD}_{600}$  (triangle)

Table 2-  $\text{NH}_4^+\text{-N}$ ,  $\text{NO}_2^-\text{-N}$ , and  $\text{NO}_3^-\text{-N}$  Removal Efficiency and Rate by *A. calcoaceticus* SCC2, *P. monteilii* Nht, *P. mendocina* AquaN, *R. erythropolis* R1 and Their Consortium after 9 h, 9 h, and 36 h Incubation

| Strain                       | $\text{NH}_4^+\text{-N}$ removal efficiency (%) | $\text{NH}_4^+\text{-N}$ removal rate ( $\text{mg.L}^{-1}.\text{h}^{-1}$ ) | $\text{NO}_2^-\text{-N}$ removal efficiency (%) | $\text{NO}_2^-\text{-N}$ removal rate ( $\text{mg.L}^{-1}.\text{h}^{-1}$ ) | $\text{NO}_3^-\text{-N}$ removal efficiency (%) | $\text{NO}_3^-\text{-N}$ removal rate ( $\text{mg.L}^{-1}.\text{h}^{-1}$ ) |
|------------------------------|---|--|---|--|---|--|
| <i>A. calcoaceticus</i> SCC2 | $99.3 \pm 0.20^a$                               | $13.23 \pm 0.74^a$   | $98.0 \pm 0.45^a$                               | $1.38 \pm 0.38^b$  | $21.2 \pm 2.6^c$                                | $0.67 \pm 0.12^d$  |
| <i>P. monteilii</i> Nht,     | $95.0 \pm 0.05^a$                               | $10.92 \pm 0.43^b$   | $97.2 \pm 2.10^a$                               | $1.1 \pm 0.03^c$   | $50.9 \pm 3.52^a$                               | $0.84 \pm 0.13^c$  |
| <i>P. mendocina</i> AquaN    | $82.3 \pm 1.30^b$                               | $7.62 \pm 1.80^c$  | $96.8 \pm 0.20^a$                               | $1.1 \pm 0.00^c$   | $47.4 \pm 6.27^a$                               | $1.76 \pm 0.20^a$  |
| <i>R. erythropolis</i> R1    | $19.3 \pm 3.75^c$                               | $1.86 \pm 1.01^d$  | $0.0^c$   | $0.00^d$   | $3.8 \pm 1.17^d$                                | $0.1 \pm 0.03^b$   |
| Bacterial consortium         | $98.3 \pm 0.36^a$                               | $9.35 \pm 0.85^{bc}$   | $90.5 \pm 2.20^b$                               | $2.42 \pm 0.44^a$  | $36.7 \pm 7.11^b$                               | $1.03 \pm 0.21^b$  |

The bacterial consortium reduced 102.5 mg.L<sup>-1</sup> of NO<sub>3</sub><sup>-</sup>-N during the first 12 hours and then the nitrate concentration remained constant. Just after nitrate-reducing, nitrite was increased to 4.0 mg.L<sup>-1</sup>. The Aerobic denitrification removal efficiency and rate in the presence of nitrite and nitrate for pure cultures and consortium are shown in Table 3.

The resting cells of the bacterial consortium removed ammonium within 12 h time as shown in Figure 6. Nitrite and nitrate were not detected in the resting cell supernatants. Total nitrogen analysis detected 65.9% nitrogen removal during the resting cells assay.

**High-strength Ammonium Removal by the Consortium:** The high-strength NH<sub>4</sub><sup>+</sup>-N removal efficiency of the bacterial consortium under different NH<sub>4</sub><sup>+</sup>-N loads was shown in Table 4. Consortium removed completely 249 mg.L<sup>-1</sup> NH<sub>4</sub><sup>+</sup>-N within 24 h, but subsequent higher concentrations were degraded by more than 50% efficiency at 24 h and approximately 100% efficiency after 48 h. These results showed consortium can completely degrade 2045 mg.L<sup>-1</sup> of NH<sub>4</sub><sup>+</sup>-N during 48 h. The NH<sub>4</sub><sup>+</sup>-N removal performance at the concentration of 2045 mg.L<sup>-1</sup> by bacterial consortium showed a statistically significant difference compared to pure cultures (Fig. 7).

Table 3- Removal Efficiency of High Strength Ammonium after 24 h and 48 h Incubation by Bacterial Consortium

| NH <sub>4</sub> <sup>+</sup> -N Concentration (mg.L <sup>-1</sup> ) | NH <sub>4</sub> <sup>+</sup> -N removal efficiency (%) after 24 h | NH <sub>4</sub> <sup>+</sup> -N removal efficiency (%) after 48 h |
|---|---|---|
| 249   | 100 ± 0.00  | 100 ± 0.00  |
| 426   | 63.1 ± 11.24  | 99.7 ± 0.00   |
| 575   | 54.3 ± 11.87  | 99.2 ± 0.21   |
| 743   | 65.4 ± 2.12   | 99.5 ± 0.14   |
| 990   | 53.1 ± 4.31   | 99.8 ± 0.07   |
| 1155  | 57.2 ± 5.51   | 99.8 ± 0.14   |
| 1648  | N.D*  | 99.9 ± 0.00   |
| 2045  | N.D   | 99.9 ± 0.00   |

\* Not Detected




Fig. 6- Ammonium removal characteristics by resting cells of the bacterial consortium at 30 °C in HNM. a: Ammonium removal characteristics, b: Total nitrogen removal characteristics. Error bars indicate the standard deviation of the mean for three replicates. Figure characterizes NH<sub>4</sub><sup>+</sup>-N (diamond), NO<sub>2</sub><sup>-</sup>-N (circle), and NO<sub>3</sub><sup>-</sup>-N (square)

Table 4- Removal Efficiency of High Strength Ammonium (2045 mg.L<sup>-1</sup>) after 48 h Incubation by Pure Cultures and Bacterial Consortium

| Strain                | NH <sub>4</sub> <sup>+</sup> -N removal efficiency (%) |
|-----------------------|--|
| A. calcoaceticus SCC2 | 47.1 ± 2.3a  |
| P. monteilii Nht,     | 77.6 ± 1.2b  |
| P. mendocina AquaN    | 79.6 ± 0.2b  |
| R. erythropolis R1    | 74.7 ± 2.4c  |
| Bacterial consortium  | 99.9 ± 0.0d  |

Data in the same column with different alphabetical superscripts differ significantly by one-way ANOVA with Duncan's test (p<0.05)




Fig. 7- Characteristics of high strength NH<sub>4</sub><sup>+</sup>-N (2045 mg.L<sup>-1</sup>) removal of the bacterial consortium and pure cultures at 30 °C after 48 h incubation. The inoculum size was 5% v/v. Error bars indicate the standard deviation of the mean for three replicates. Different alphabetical superscripts indicate significant differences by one-way ANOVA with Duncan's test (p < 0.05)

## Discussion and Conclusion

Currently, heterotrophic nitrification-aerobic denitrification bacteria attract researchers' attention because they are able to degrade ammonium, nitrite, and nitrate in aerobic conditions and are suitable to use in the bioremediation of water and wastewater (31).

In this study, the HN-AD bacteria were selected and their consortium was designed. Two strains *Pseudomonas monteilii* Nht (99.71% similarity) and *Pseudomonas mendocina* AquaN (99.71% similarity) were isolated from soil and aquarium water, respectively. The bacterial consortium consisted of *A. calcoaceticus* SCC2, *R. erythropolis* R1, *P. mendocina* AquaN, and *P. monteilii* Nht. The interaction experiments did not show any interference between the strains which this result indicated strains were suitable for consortium preparation. All strains have well performance in the HN-AD process. The single strains and consortium were able to remove ammonium, nitrite, and nitrate. *P. mendocina* AquaN, *A. calcoaceticus* SCC2, and *P. monteilii* Nht performed well in ammonium removal by up to 98.5% efficiency within 24 h; however, the performances of other strains like *Alcaligenes* sp. TB and *Enterobacter cloacae* CF-S27 were slower as well as removed ammonium by 99.42% and 81% efficiency during 44 h and 36 h, respectively (13, 20). *R. erythropolis* R1 was a slow-growing bacterium and removed 85.0% of ammonium during 24 h incubation. During denitrification in DM-1, nitrite converted to nitrate but nitrate was not removed. Also, the result of denitrification in DM-2 showed pure cultures consumed a part of nitrate. Similar results were also reported in previous studies such as *Aeromonas* sp. HN-02 and *Arthrobacter nicotianae* D51 removed 33.32% and 65% of nitrate in aerobic conditions, respectively (32, 33).

According to Stein (34), aerobic denitrifying bacteria reduced nitrate and oxygen simultaneously to maximize respiration, and due to this process nitrate was not completely removed in denitrification tests (35). It can be concluded that the strains have little ability in nitrate denitrification and the nitrate reduction was either assimilated by the cells or consumed in competitive denitrification with oxygen. Many studies have indicated that HN-AD bacteria have different complex nitrogen metabolic pathways, which transform inorganic-N into organic or gaseous nitrogen by partial nitrification, denitrification, assimilation, and dissimilation (16). The pure cultures did not produce any nitrite and nitrate during the nitrification process because of their HN-AD pathways. Based on the studies of the HN-AD genes, the HN-AD pathway can be as follows: ammonium could convert to nitrite and then to nitrate. Afterward, nitrate rapidly entered aerobic denitrification processes and was removed. In some cases, nitrite was used directly in aerobic denitrification (13, 33). In some cases, ammonium was converted to nitrogen gaseous without being converted to nitrite and nitrate (36).

Remarkably, microbial communications stabilize their cooperative behaviors and improve enzyme activity and biomass yield (8, 37, 38). Therefore, bacteria in the consortium can complete each nitrogen metabolic pathway and accelerate activities. In this research, the bacterial consortium also showed a significantly higher  $\text{NH}_4^+$ -N removal efficiency (98.3%) than their pure cultures at the end of 9 h cultivation, nitrite and nitrate were not produced at this time which is similar to the pure culture's performances. These results investigated enhancement enzyme function in the consortium state. In Angar, Kebbouche-Gana (37), the consortium of *Alcaligenes* sp. S84S3 and *Proteus* sp. S19

removed 65% of the ammonium as high as 272.72 mg.L<sup>-1</sup> NH<sub>4</sub><sup>+</sup>-N during 69 h incubation under optimal conditions. In another study, the consortium of *Pseudomonas fulva* K3, *Pseudomonas mosselii* K17, and *Enterobacter hormaechei* A16 removed 99.62% of the ammonium as high as 100 mg.L<sup>-1</sup> NH<sub>4</sub><sup>+</sup>-N during 12 h incubation that was significantly better than pure cultures removal function (39).

Ammonium removal analysis of consortium resting cells showed increasing in NH<sub>4</sub><sup>+</sup>-N concentration at first 2 h due to the activation of proteolytic enzymes at the end of the logarithmic growth phase and the release of the dead bacterial cells into the medium (32, 40). At the end of the resting cell test, 65.9% of total nitrogen was removed from the medium; this value was expected to be converted to gaseous. This means that the bacterial consortium had nitrification ability after passing the growth phase and in a high-density condition. The important result of this study is a higher tolerance of consortium to ammonium and a higher biodegradation rate than previous results reported by other co-cultures. The consortium tolerated and grew in a medium with 250 to 2045 mg.L<sup>-1</sup> NH<sub>4</sub><sup>+</sup>-N concentration and degraded up to 95% of NH<sub>4</sub><sup>+</sup>-N during 48 h, with significantly high degradation efficiency than the pure culture of bacteria. However, the high-strength ammonium degradation reported earlier. *Pseudomonas stutzeri* GEP-01 removed 1000 mg.L<sup>-1</sup> NH<sub>4</sub><sup>+</sup>-N by 78.9% efficiency within 72 h (41). In addition, *Alcaligenes faecalis* TF-1 tolerated 1963.94 mg.L<sup>-1</sup> NH<sub>4</sub><sup>+</sup>-N and removed by 65% efficiency after 144 h (36).

The bacterial consortium had a denitrification function in the presence of 23 mg.L<sup>-1</sup> NO<sub>2</sub><sup>-</sup>-N. This concentration was higher than the tolerable concentration for the pure cultures. The bacterial consortium removed 40% of NO<sub>3</sub><sup>-</sup>-N in DM-2 at first 12

h. This result was in contrast to pure culture performances because NO<sub>3</sub><sup>-</sup>-N concentration in the consortium test did not fluctuate and remained stable as well as pure cultures started to consume nitrate after 12 h.

In this study, HN-AD strains including *P. mendocina* AquaN and *P. monteilii* Nht were isolated and designed for the consortium with *A. calcoaceticus* SCC2, and *R. erythropolis* R1. These strains could utilize ammonium, nitrate, and nitrite as the sole nitrogen sources. The bacterial consortium performed better HN-AD function than single strains. In addition, the consortium exhibited high tolerance capacity on 2045 mg.L<sup>-1</sup> NH<sub>4</sub><sup>+</sup>-N and was able to remove such concentrated ammonium completely within 48 h. Thus, the co-culture and interactions between HN-AD strains also contribute to high-efficiency ammonium removal.

### Acknowledgments

This study is part of a master's thesis that has been financially supported by the University of Isfahan, Iran. We also thank Fatemeh Ghoreishi for providing *Acinetobacter* for this research.

### Declaration of interest statement

The authors declare that they have no conflict of interest.

### Data availability statement

All data generated or analyzed during this study are included in this published article. "Sequence" data that support the part of the findings of this study have been deposited in "GenBank" with the accession codes MN167837.1 and MN435149.1.

### References

- (1) Fields S. Global nitrogen: Cycling out of control. *Environmental Health Perspectives* 2004; 112 (10): 557-63. doi: 10.1289/ehp.112-a556

- (2) Wang X., Wang W., Zhang Y., Sun Z., Zhang J., Chen G., Li J. Simultaneous nitrification and denitrification by a novel isolated *Pseudomonas* sp. JQ-H3 using polycaprolactone as carbon source. *Bioresource Technology* 2019; 288: 121506. doi: 10.1016/j.biortech.2019.121506
- (3) Trung Tran T., Bott NJ., Dai Lam N., Trung Nguyen N., Hoang Thi Dang O., Hoang Le D., Tung Le L., Hoang Chu H. The role of *Pseudomonas* in heterotrophic nitrification: A case study on shrimp ponds (*Litopenaeus vannamei*) in Soc Trang province. *Microorganisms* 2019; 7 (6): 155. doi: 10.3390/microorganisms7060155
- (4) Liu H., Chen Z., Guan Y., Xu S. Role and application of iron in water treatment for nitrogen removal: A review. *Chemosphere* 2018; 204: 51-62. doi: 10.1016/j.chemosphere.2018.04.019
- (5) Boroumand Y., Razmjou A., Moazzam P., Mohagheghian F., Eshaghi G., Etemadifar Z., Asadnia M., Shafiei R. Mussel inspired bacterial denitrification of water using fractal patterns of polydopamine. *Journal of Water Process Engineering* 2020; 33: 101105. doi: 10.1016/j.jwpe.2019.101105
- (6) Mook WT., Chakrabarti MH., Aroua MK., Khan GM., Ali BS., Islam MS., Hassan MA. Removal of total ammonia nitrogen (TAN), nitrate and total organic carbon (TOC) from aquaculture wastewater using electrochemical technology: A review. *Desalination* 2012; 285: 1-13. doi: 10.1016/j.desal.2011.09.029
- (7) Huang F., Pan L., He Z., Zhang M., Zhang M. Culturable heterotrophic nitrification-aerobic denitrification bacterial consortia with cooperative interactions for removing ammonia and nitrite nitrogen in mariculture effluents. *Aquaculture* 2020; 523: 735211. doi: 10.1016/j.aquaculture.2020.735211
- (8) Zhang J., Wu P., Hao B., Yu Z. Heterotrophic nitrification and aerobic denitrification by the bacterium *Pseudomonas stutzeri* YZN-001. *Bioresource Technology* 2011; 102 (21): 9866-9. doi: 10.1016/j.biortech.2011.07.118
- (9) Lu H., Weng Z., Wei H., Zhou J., Wang J., Liu G., Guo W. Simultaneous bisphenol F degradation, heterotrophic nitrification and aerobic denitrification by a bacterial consortium. *Journal of Chemical Technology & Biotechnology* 2017; 92 (4): 854-60. doi: 10.1002/jctb.5069
- (10) Xi H., Zhou X., Arslan M., Luo Z., Wei J., Wu Z., El-Din MG. Heterotrophic nitrification and aerobic denitrification process: Promising but a long way to go in the wastewater treatment. *Science of The Total Environment* 2022; 805: 150212. doi: 10.1016/j.scitotenv.2021.150212
- (11) Sarioglu OF., Suluyayla R., Tekinay T. Heterotrophic ammonium removal by a novel hatchery isolate *Acinetobacter calcoaceticus* STB1. *International Biodeterioration & Biodegradation* 2012; 71: 67-71. doi: 10.1016/j.ibiod.2012.04.012
- (12) Chen J., Gu S., Hao H., Chen J. Characteristics and metabolic pathway of *Alcaligenes* sp. TB for simultaneous heterotrophic nitrification-aerobic denitrification. *Applied Microbiology and Biotechnology* 2016; 100 (22): 9787-94. doi: 10.1007/s00253-016-7840-x
- (13) Padhi SK., Tripathy S., Sen R., Mahapatra AS., Mohanty S., Maiti NK. Characterisation of heterotrophic nitrifying and aerobic denitrifying *Klebsiella pneumoniae* CF-S9 strain for bioremediation of wastewater. *International Biodeterioration & Biodegradation* 2013; 78: 67-73. doi: 10.1016/j.ibiod.2013.01.001
- (14) Kim JK., Park KJ., Cho KS., Nam SW., Park TJ., Bajpai R. Aerobic nitrification-denitrification by heterotrophic *Bacillus* strains. *Bioresource Technology* 2005; 96 (17): 1897-906. doi: 10.1016/j.biortech.2005.01.040
- (15) Gao J., Zhu T., Liu C., Zhang J., Gao J., Zhang J., Cai M., Li Y. Ammonium removal characteristics of heterotrophic nitrifying bacterium *Pseudomonas stutzeri* GEP-01 with potential for treatment of ammonium-rich wastewater. *Bioprocess and Biosystems Engineering* 2020; 43: 959-69. doi: 10.1007/s00449-020-02292-x
- (16) Jin P., Chen Y., Yao R., Zheng Z., Du Q. New insight into the nitrogen metabolism of

- simultaneous heterotrophic nitrification-aerobic denitrification bacterium in mRNA expression. *Journal of Hazardous Materials* 2019; 371: 295-303. doi:10.1016/j.jhazmat.2019.03.023
- (17) Bhatia SK., Bhatia RK., Choi YK., Kan E., Kim YG., Yang YH. Biotechnological potential of microbial consortia and future perspectives. *Critical Reviews in Biotechnology* 2018; 38 (8): 1209-29. doi: 10.1080/07388551.2018.1471445
- (18) Derikvand P., Etemadifar Z., Biria D. Taguchi optimization of dibenzothiophene biodesulfurization by *Rhodococcus erythropolis* R1 immobilized cells in a biphasic system. *International Biodeterioration & Biodegradation* 2014; 86: 343-8. doi: 10.1016/j.ibiod.2013.10.006
- (19) Ghoreishi F., Etemadifar Z. Heavy metal removal by phosphate solubilizing *Acinetobacter calcoaceticus* isolated from rhizosphere. *Journal of Biology* 2017; 6 (11): 230-9. doi: 10.15412/J.JBTW.01061104
- (20) Zhang QL., Liu Y., Ai GM., Miao LL., Zheng HY., Liu ZP. The characteristics of a novel heterotrophic nitrification-aerobic denitrification bacterium, *Bacillus methylotrophicus* strain L7. *Bioresource Technology* 2012; 108: 35-44. doi: 10.1016/j.biortech.2011.12.139
- (21) Falk S., Liu B., Braker G. Isolation, genetic and functional characterization of novel soil nirK-type denitrifiers. *Systematic and Applied Microbiology* 2010; 33 (6): 337-47. doi: 10.1016/j.syapm.2010.06.004
- (22) Brenner DJ., Krieg NR., Staley JT. *Bergey's manual of systematics of Archaea and Bacteria*. 2nd ed. Michigan: Springer; 2015.
- (23) Wang L., Fu Y., Wang S., Ye F., Cui E., Sun Q. Domesticating a halotolerant bacterium of vibrio sp. LY1024 with Heterotrophic nitrification-aerobic denitrification property for efficient Nitrogen removal in mariculture wastewater treatment. *Coatings* 2022; 12 (11): 1786. doi: 10.3390/coatings12111786
- (24) Premalatha A., Karthi A. Phytochemical screening and antimicrobial activity of leaf of *Atalantia monophylla* (L.) Corr (rutaceae). *International Journal of Pharmacy and Biological Sciences* 2017; 7 (4): 179-84.
- (25) Slama HB., Cherif-Silini H., Chenari Bouket A., Qader M., Silini A., Yahiaoui B., Alenezi FN., Luptakova L., Triki MA., Vallat A., Oszako T. Screening for *Fusarium* antagonistic bacteria from contrasting niches designated the endophyte *Bacillus halotolerans* as plant warden against *Fusarium*. *Frontiers in Microbiology* 2019; 9: 3236. doi: 10.3389/fmicb.2018.03236
- (26) APHA. *Standard methods for the examination of water and wastewater*. 20th ed. Washington, DC: American Public Health Association; 1999.
- (27) Daniels L., Hanson RS., Phillips JA. Chemical analysis. In: Reddy CA, Beveridge TJ, Breznak JA, Marzluf GA, Schmidt TM, Snyder LR, editors. *Methods for General and Molecular Microbiology*. 3rd ed. Washington, D.C: American Society of Microbiology; 2007: 462-503.
- (28) Giustarini D., Rossi R., Milzani A., Dalle-Donne I. Nitrite and nitrate measurement by Griess reagent in human plasma: Evaluation of interferences and standardization. *Methods in enzymology* 2008; 440: 361-80. doi: 10.1016/S0076-6879(07)00823-3
- (29) Feng Y., Feng J., Shu Q. Isolation and characterization of heterotrophic nitrifying and aerobic denitrifying *Klebsiella pneumoniae* and *Klebsiella variicola* strains from various environments. *Journal of Applied Microbiology* 2018; 124 (5): 1195-211. doi: 10.1111/jam.13703
- (30) Ramis-Ramos G. ANTIOXIDANTS Synthetic Antioxidants. In: Caballero B, editor. *Encyclopedia of Food Sciences and Nutrition*. 2nd ed. Burjassot: Elsevier; 2003: 265-75.
- (31) Padhi SK., Tripathy S., Mohanty S., Maiti NK. Aerobic and heterotrophic nitrogen removal by *Enterobacter cloacae* CF-S27 with efficient utilization of hydroxylamine. *Bioresource Technology* 2017; 232: 285-96. doi: 10.1016/j.biortech.2017.02.049
- (32) Cai X., Li K., He T., Wang Y., Zhang X.,

- Xie E., Ding N., Li Z. Characteristics of heterotrophic nitrifying and aerobic denitrifying *Arthrobacter nicotianae* D51 strain in the presence of copper. *Water* 2019; 11 (3): 434. doi: 10.3390/w11030434
- (33) He X., Sun Q., Xu T., Dai M., Wei D. Removal of nitrogen by heterotrophic nitrification–aerobic denitrification of a novel halotolerant bacterium *Pseudomonas mendocina* TJPU04. *Bioprocess and Biosystems Engineering* 2019; 42 (5): 853-66. doi: 10.1007/s00449-019-02088-8
- (34) Stein LY. Heterotrophic nitrification and nitrifier denitrification. In: Ward BBA, Daniel J. Klotz, Martin G, editors. *Nitrification*. New Jersey: Wiley Online Library; 2011: 95-114.
- (35) Yang L., Wang XH., Cui S., Ren YX., Yu J., Chen N., Xiao Q., Guo LK., Wang RH. Simultaneous removal of nitrogen and phosphorous by heterotrophic nitrification-aerobic denitrification of a metal resistant bacterium *Pseudomonas putida* strain NP5. *Bioresource Technology* 2019; 285: 121360. doi: 10.1016/j.biortech.2019.121360
- (36) Zhang Q., Zhu Y., Yuan C., Zhang C., Cui M., Zhao T. Nitrogen removal and mechanism of an extremely high-ammonia tolerant heterotrophic nitrification-aerobic denitrification bacterium *Alcaligenes faecalis* TF-1. *Bioresource Technology* 2022; 361: 127643. doi: 10.1016/j.biortech.2022.127643
- (37) Angar Y., Kebbouche-Gana S., Djelali NE., Khemili-Talbi S. Novel approach for the ammonium removal by simultaneous heterotrophic nitrification and denitrification using a novel bacterial species co-culture. *World Journal of Microbiology and Biotechnology* 2016; 32 (3): 36. doi: 10.1007/s11274-015-2007-y
- (38) Luo G., Zhang N., Cai S., Tan H., Liu Z. Nitrogen dynamics, bacterial community composition and biofloc quality in biofloc-based systems cultured *Oreochromis niloticus* with poly- $\beta$ -hydroxybutyric and polycaprolactone as external carbohydrates. *Aquaculture* 2017; 479: 732-41. doi: 10.1016/j.aquaculture.2017.07.017
- (39) Hu J., Su Q., Xiao C., Deng X., Liu X., Feng J., Chi R. Removal of ammonia nitrogen from residual ammonium leaching solution by heterotrophic nitrification-aerobic denitrification process. *Environmental Technology* 2022; 1-12. doi: 10.1080/09593330.2022.2064235
- (40) Carroll KC., Morse SA., Mietzner TA., Miller S., Jawetz, M. *Adelberg's medical microbiology*. 27th ed. Stamford: Lange Medical Books/McGraw-Hill, Medical Pub. Division; 2015.
- (41) Sukeetha DS., Kumar SD., Manikandan A., Akila V., Santhanam P., Rajakumar S. Effect of immobilized nitrifying bacterial consortium on ammonia biodegradation in aquaculture pond and enhanced growth of *Labeo rohita*: An in vitro and in vivo studies. *Arabian Journal for Science and Engineering* 2020; 45 (1): 1-13. doi: 10.1007/s13369-019-04073-5